

# Colour ratios of pulsation amplitudes in red giants

J.P. Huber, T.R. Bedding & S.J. O'Toole

*School of Physics, University of Sydney 2006, Australia*

**Abstract.** We have analysed the three-colour photometry of 34 pulsating red giants that was published by Percy, Wilson and Henry (2001). As is common with semiregulars, many of these stars have long secondary periods (LSPs) of several years, in addition to the usual shorter-period pulsations. We measured pulsation amplitudes in  $VRI$  and found that both the  $V/R$  and  $V/I$  amplitude ratios in most stars are systematically lower for the LSPs than for the shorter periods. These results must be taken into account by any theory that attempts to explain LSPs.

**Keywords:** stars: AGB and post-AGB – stars: oscillations

## 1. Introduction

Many semiregular variables have long secondary periods (LSPs) of several years, in addition to their shorter-period pulsations (Houk, 1963; Kiss et al., 1999). The origin of LSPs is unclear, and may be due to binarity (Wood et al., 1999; Wood et al., 2001) or to strange pulsation modes caused by convection-pulsation interaction (Wood, 2000). Multi-colour photometry may shed some light on this puzzle.

## 2. Data Analysis

Percy et al. (2001) have published three-colour ( $VRI$ ) photometry of 34 pulsating red giants covering more than ten years. These data are publicly available. We have calculated amplitude spectra and, for stars showing more than one period, we measured the amplitudes of the two highest peaks. This was done in each of the three wavebands, allowing us to calculate amplitude ratios.

## 3. Results

Figures 1 and 2 show the  $V/R$  and  $V/I$  amplitude ratios for all stars with two periods. In most cases, the longer period is an LSP, with a value of several hundred days. The figures show that in a given star, the amplitude of the LSP generally has less wavelength dependence than does the shorter period. This result must be taken into account by any



© 2008 Kluwer Academic Publishers. Printed in the Netherlands.

Table I. Spectral types and periods for a subset of the stars

| Name           | Spectral Type | Periods (days) |
|----------------|---------------|----------------|
| 30 Her (g Her) | M6 III        | 871            |
| 4 Ori          | S3.5/1        | 386, 35        |
| BC CMi         | M4 III        | 327, 38        |
| EG And         | M2e           | 233, 28        |
| $\mu$ Gem      | M3 III        | 2547, 26       |
| UW Lyn         | M3 IIIab      | 506, 37        |
| V1070 Cyg      | M7 III        | 486, 43        |
| V642 Her       | M4 III        | 2547, 25       |

theory that attempts to explain LSPs. Note that EG And is known to be a symbiotic binary, and the LSP corresponds to the orbital period. Interestingly, the amplitude ratios of EG And are similar to those of other stars, perhaps lending some weight to the binary model as the explanation of LSPs.

#### 4. Future Study

Further analysis, perhaps using Period 98, should give more accurate amplitude ratios. Comparison with observed velocities may help determine which stars are members of a binary system (Wood et al., 2001; Hinkle et al., 2002).

#### References

- Hinkle K. H., Lebzelter T., Joyce R. R., Fekel F. C., 2002, *AJ*, 123, 1002  
Houk, N., 1963, *AJ*, 68, 253  
Kiss, L. L., Szatmáry, K., Cadmus, R. R., & Mattei, J. A., 1999, *A&A*, 346, 542.  
Percy, J.R., Wilson, J.B., & Henry, G. W., 2001, *PASP*, 113, 983  
Wood, P.R., 2000, *Proc. Astron. Soc. Aust.*, 17, 18  
Wood, P. R., Alcock, C., Allsman, R. A., et al., 1999, in *IAU Symp. 191: Asymptotic Giant Branch Stars*, eds. T. Le Bertre, A. Lebre & C. Waelkens, p. 151  
Wood, P. R., Axelrod, T. S., & Welch, D. L. 2001, *ASP Conf. Ser. 229: Evolution of Binary and Multiple Star Systems*, eds. Ph. Podsiadlowski, S. Rappaport, A. R. King, F. D'Antona and L. Burder, p. 233




Figure 1. Amplitude ratios  $V/R$  and  $V/I$  for stars with two periods




Figure 2. Same as Fig. 1, but restricted to stars classified as M giants